

PP1495 (Revised)

Management of Rhizoctonia Root and Crown Rot of Sugar Beets

Reviewed by

Mohamed F.R. Khan, Extension Sugarbeet Specialist, Department of Plant Pathology
North Dakota State University, Fargo and University of Minnesota, St. Paul

Melvin D. Bolton, U.S. Department of Agriculture,
Agricultural Research Service, Northern Crop Science Laboratory, Fargo

Rhizoctonia root and crown rot is one of the most severe soil-borne diseases of sugar beets and it's a major problem for growers in Minnesota and North Dakota.

The local sugar beet-growing areas have been in a wet cycle during the past 20 years. Fields with high disease severity probably have resulted from a combination of wet conditions and a prevalence of other susceptible host crops in the rotation. For example, nonsusceptible crops such as wheat and barley formerly used in rotation with sugar beets have decreased in acreage and have been replaced with susceptible crops such as soybeans, edible beans and corn.

Rhizoctonia root and crown rot may reduce yield significantly and diseased beets may cause problems in storage piles. Growers at some sugar cooperatives are required to destroy fields with more than a 50 percent incidence of root rot because infected roots cannot be placed in piles for long term storage.

Causal agent

Rhizoctonia root and crown rot is caused by the fungus *Rhizoctonia solani*. This fungus is divided into a number of anastomosis groups (AGs) based on vegetative compatibility reactions that occur when hyphae of two similar isolates fuse and genetic material is exchanged.

The main AGs that seriously impact sugar beets in Minnesota and North Dakota are AG 2-2 IV and AG 2-2 IIIB. The pathogen is present in all soils but becomes problematic in fields where the pathogen population is high because of the frequent use of susceptible host crops in the rotation.



Figure 1. Wilting of oldest leaves first and yellowing of plants infected with *R. solani*. (NDSU photo)



Figure 2. Blackening of petioles and infection at the soil level. (NDSU photo)

NDSU EXTENSION
SERVICE

North Dakota State University, Fargo, North Dakota

Revised September 2016



Disease symptoms

The most common symptom is wilting of the leaves, starting with the oldest leaves, and is likely a consequence of damage to the vascular system (Figures 1 and 3). Leaves may or may not become yellow. In some instances, the petioles may develop some blackening where they are attached to the crown (Figure 2).

Crown rot infection probably is initiated when infected soil is thrown into crowns during cultivation, by rain splash of infected soil into the crowns or where the petiole attachment to the crown is covered with infected soil (Figure 2). Root rot infection typically starts at or just below the soil line (Figure 3), and sometimes lower on the tap root (Figure 4).

Root rot symptoms may range from scattered brown to black lesions on the root surface to complete rotting of the root. Initially, the disease may occur in a few patches in a field. If significant inoculum is present, often as a result of continuous planting of susceptible hosts, entire sugar beet fields may be lost to the disease when conditions are favorable for disease development.

Disease management

Rotations with crops such as wheat and barley, which are not hosts of *R. solani* AG 2-2 IIIB or AG 2-2 IV, will help reduce the inoculum pressure. Because the pathogen is more severe in wet conditions, draining and leveling of fields will help in disease management.

Fields with a history of *Rhizoctonia* root and crown rot should be planted to a variety with good disease resistance. The fungicides azoxystrobin and prothioconazole with a non-ionic surfactant at 0.125 percent volume per volume (v/v), and a mixture of pyraclostrobin and fluxapyroxad applied in a 7-inch band before infection takes place provide effective disease control.

Fungicides should be applied when the average daily soil temperature at the 4-inch soil depth is about 60 to 62 F or before the canopy covers the rows (four- to 10-leaf stage).

Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply endorsement.



Figure 3. *Rhizoctonia* root rot infected plant with yellow leaves, and showing infection start on the root just below the soil line. (NDSU photo)



Figure 4. *Rhizoctonia solani*-infected plant with rot symptoms on the lower part of the tap root. (NDSU photo)

The NDSU Extension Service does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names. NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

For more information on this and other topics, see www.ag.ndsu.edu

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. North Dakota State University does not discriminate on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, physical and mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, or status as a U.S. veteran. Direct inquiries to: Vice Provost for Faculty and Equity and Title IX/ADA Coordinator, Old Main 201, 701-231-7708. This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881.