







Figure 2. Relationship between Rhizoctonia root rot rating and amount of DNA of *Rhizoctonia solani* AG 2-2 in soil.

#### ACKNOWLEDGEMENTS

We thank the Sugarbeet Research and Education Board of Minnesota and North Dakota for funding in support of this research; the University of Minnesota, Northwest Research and Outreach Center for providing facilities and equipment; Katie Sheetz and Claire Carlson for technical assistance; and agriculturists from sugarbeet cooperatives for locating fields with a history of Rhizoctonia.

#### LITERATURE CITED

1. Brantner, J.R. 2015. Plant pathology laboratory: summary of 2013-2014 field samples. 2014 Sugarbeet Res. Ext. Rept. 44:138-139.
2. Brantner, J.R. and C.E. Windels. 2011. Plant pathology laboratory: summary of 2009-2010 field samples. 2010 Sugarbeet Res. Ext. Rept. 41:260-261.
3. Brantner, J.R. and C.E. Windels. 2009. Plant pathology laboratory: summary of 2007-2008 field samples. 2008 Sugarbeet Res. Ext. Rept. 39:250-251.
4. Budge, G.E., Shaw, M.W., Colyer, A., Pietravalle, S., and N. Boonham. 2009. Molecular tools to investigate *Rhizoctonia solani* distribution in soil. Plant Pathol. 58:1071-1080.
5. Chanda, A. K., Brantner, J. R., Metzger, M., Bloomquist, M., and Groen, C. 2017. Integrated Management of Rhizoctonia on Sugarbeet with Varietal Resistance, At-Planting Treatments and Postemergence Fungicides. 2016 Sugarbeet Res. Ext. Rept. 47:174-179.
6. Crane, E., Brantner, J.R., and C.E. Windels. 2013. Plant pathology laboratory: summary of 2011-2012 field samples. 2012 Sugarbeet Res. Ext. Rept. 43:169-170.